

# THE BRAND MOMS TURN TO FIRST\*

\*When they first treat their children's upper respiratory allergies.



FREE OF  
DYE, SUGAR,  
& ALCOHOL

**Children's  
BENADRYL®  
Allergy Liquid†**

Use products only as directed.



**Children's  
BENADRYL®  
Allergy Plus  
Congestion†‡**

Use product only as directed.



**Children's  
BENADRYL®  
Chewables†**

Use product only as directed.

AGE	DOSE
<b>6-11 years</b>	Take 5 to 10 mL every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
<b>2-5 years</b>	Do not use unless directed by a doctor.
<b>Under 2 years</b>	Do not use.

AGE	DOSE
<b>12+ years</b>	Take 10 mL every 4 hours. Do not take more than 6 doses in 24 hours.
<b>6-11 years</b>	Take 5 mL every 4 hours. Do not take more than 6 doses in 24 hours.
<b>4-5 years</b>	Do not use unless directed by a doctor.
<b>Under 4 years</b>	Do not use.

AGE	DOSE
<b>12+ years</b>	2 to 4 chewable tablets (25 mg to 50 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
<b>6-11 years</b>	1 to 2 chewable tablets (12.5 mg to 25 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
<b>2-5 years</b>	Do not use unless directed by a doctor.
<b>Under 2 years</b>	Do not use.



Help your child find relief from most outdoor itches with **BENADRYL® Extra Strength Itch Stopping Cream.**

**For additional information on allergy management, talk to your pediatrician.**

†Relieves runny nose, sneezing, itchy, watery eyes, itching of the nose or throat.

‡Use only with dosing cup provided with product.

mL = milliliter